



# Pheromone Chemicals

The name you can always trust

Mfrs: Pheromone Traps, Lures, Fruit Fly Traps, Sticky traps

## *Earias insulana* (Spiny bollworm)

The spiny bollworm (*Earias insulana*), is the most important pest of Cotton and Okra.

**Host range:** Okra, Cotton, Aibica, Hollyhock (*A. rosea.*), Safflower (*Carthamus sp.*), Indian mallow (*Abutilon sp.*), Corchorus sp., Hibiscus sp., Malvas sp., Malvastrum sp., Sida sp., Theobroma sp. and Urena sp



### Nature of Damage

The caterpillars of the pest not only bore the bolls but also attack the shoots, buds and flowers. The infested bolls open prematurely and produce poor lint resulting in lower market value.

Use Pheromone Traps from 1 month crop stage @ 6 – 8 No's per acre to control pest at early stage.

Trap canopy should be placed one foot above crop canopy to achieve optimum catch.

Always use Phero – Sensor™ – SP / BP for best results.

**Pc** Pheromone Lures for  
**EARIAS INSULANA**  
Host: Cotton, Okra

Identify the pest incidence early. Stop egg laying. Get good yields

TS: Plot No.23, TSIIC Techpark, Road No.15, IDA Nacharam, Hyderabad – 76, TS, INDIA , Cell: +91-94408 97918

AP: Shop No.15, Sri Ranganayaka Swamy Complex, Patnam Bazar, GUNTUR – 2, AP Cell: +91-99120 64545

E-mail: [pheromonechemicals@gmail.com](mailto:pheromonechemicals@gmail.com), Web: [www.pheromonechemicals.in](http://www.pheromonechemicals.in)

INNOVATIVE, AFFORDABLE & QUALITY PEST MONITORING AND CONTROL SYSTEMS